



**SIFLON**

Drugs & Pharmaceuticals Pvt. Ltd.

**POULTRY**

[www.siflon.com](http://www.siflon.com)

## Sulphaclozine Powder 30% w/w

# SIFCLOZINE



### Composition:

Sulphaclozine Sodium Monohydrate  
Equivalent to Sulphaclozine 30% w/w  
Excipients Q.S

**Indications:** SIFCLOZINE is indicated for treatment of coccidiosis in poultry due to infection with *Eimeria* species, Flow Typhoid due to infection with *Salmonella gallinarum* and Towl Cholera due to infection with *Pasteurella multocida*.

**Dosage:** Orally through drinking water.

### Poultry:

Coccidiosis  
1.5 to 2 gm/1Ltr of drinking water 3-5 days.

**For Bacterial Infections:** 1 gm/1 Ltr of drinking water for 3-5 days (or) as advised by the veterinarian.

For the treatment of cholera and typhoid, doses of 1 to 2 gm SIFCLOZINE per litre water for atleast 5 days are required (Or) as directed by a veterinarian.

### Composition:

Colistin Sulphate U.S.P 10%W/W  
Excipients Q.S

Colistin is a cyclic poly peptide antibiotic and has bactericidal action against selected Gram negative bacteria.

**Indications:** Poultry: Colibacillosis, Salmonellosis

### Dosage:

**Dosage & Administrations:**  
Orally through drinking water.

### Poultry:

Preventive use : 1 gram/2 liters

Treatment use : 1 gram/1 liters

(or) as advised by a veterinarian.

Note: Medicated water should be used within 24 Hours.

## Colistin Sulphate Powder 10% w/w

# SIFCOLI



### Composition:

Each gram Contains:  
Azithromycin I.P 100mg  
Equivalent to Anhydrous Azithromycin 100mg  
Excipients Q.S

**Indications:** Treatment of CCRD, Infectious Coryza and Infectious Synovitis caused by *Mycoplasma* spp

**Dosage:** Orally through drinking water.

### Poultry:

Preventive use : 1 gram/4 liters-3 days

Treatment use: 1 gram/ 1-2 liters-3-5 days. (Or) as advised by a veterinarian.

**Sailet Features:** Better absorption and distribution through oral administration. Extraordinary ability to concentrate in tissues particularly macrophages, leukocytes, fibroblast and assist in effective killing of invading organisms.

## Azithromycin Powder 10%w/w

# SIFAZIT



# SIFLON

Drugs & Pharmaceuticals Pvt. Ltd.

### Corporate Office :

Plot No. 76 & 77,  
Nagar, Madinaguda,  
Near Miyapur, Hyderabad-500 050.  
Telangana, India.

Phone : (+91-40) 23042233, 23042159

Email : [info@siflonpharma.com](mailto:info@siflonpharma.com)

[siflonpharma@rediffmail.com](mailto:siflonpharma@rediffmail.com)

### Factory :

Siflon Drugs & Pharmaceuticals Pvt. Ltd.  
Plot No. 20 & 21, ALEAP Industrial. Estate,  
Opp. JNTU Road, Pragathi Nagar,  
Hyderabad - 500 090.

Phone : +91 40 32573153

Email : [Siflonpharma@rediffmail.com](mailto:Siflonpharma@rediffmail.com)

# ENROFLOXACIN VET 20 % w/v ORAL SOLUTION

## ENROSIF-20



**Composition :**

Each ml Contains :  
Enrofloxacin IP 200 mg.

**INDICATIONS :**

For the treatment and prevention of infections & diseases caused by bacteria (Gram +ve & Gram-ve)

**ENROSIF-20** oral is the Drug of choice for the prevention and treatment Chronic Respiratory Disease (CRD) Complex, complicated by Mycoplasma spp infection .

**ENROSIF-20** oral is very effective against diarrhoea, enteritis, fowl cholera, infectious coryza.

**Dosage :**

Poultry Prevention :1 ml in 4-8 Litre of water  
Treatment 1 ml in two litre of drinking water.

# ENROFLOXACIN VET 30 % w/v ORAL SOLUTION

## ENROSIF-30

**Composition :**

Each ml Contains :  
Enrofloxacin IP 300 mg.  
Excipients q.s.

**INDICATIONS :**

For the treatment and prevention of infections & diseases caused by bacteria (Gram +ve & Gram-ve)

**ENROSIF-30** oral is the Drug of choice for the prevention and treatment Chronic Respiratory Disease (CRD) Complex, complicated by Mycoplasma spp infection .

**ENROSIF-30** oral is very effective against diarrhoea, enteritis, fowl cholera, infectious coryza.

**Dosage :**

Poultry Prevention :1 ml in 12 Liters of water  
Treatment 1 ml in 3-6 liters of drinking water.



# Levofloxacin Oral Solution 10% w/v

## LEVOSIF



**Composition:**

Each ml contains  
Levofloxacin Hemihydrate I.P  
Equivalent to Levofloxacin 100mg  
Excipients Q.S

**Indications:**

For the control and treatment of infections in poultry like CRD, CCRD, E.Coli Infections, Salmonellosis, Infectious Coryza, Dermatitis ect. Including other infections caused by Gram Positive and Gram Negative Microorganisms.

**Dosage:** Orally through drinking water

**Poultry:**

**Prevention Use:**

2-4 ml per 10 ltrs of drinking water:3-5 days

**Treatment use:**

4-8 ml per 10 ltrs of drinking water 3-5 days (or) as advised by Veterinarian

# Toltrazuril Oral Solution 2.5% w/v

## TOLTRASIF

**Composition:**

Each ml contains  
Toltrazuril 25 mg  
Excipients Q.S

**Indications:**

Coccidiosis of all stages like schizogony and gametogony stage of Eimeria spp. in chickens and turkeys.

**Dosage:** Orally through drinking water

**Poultry:** 3.5 mg Toltrazuril per kg body weight per day for 2 consecutive days, equivalent to 28ml of TOLTRASIF per 100kg of live body weight per day. The exact dosage must be carefully administered according to the live body weight and daily water consumption. TOLTRASIF should be added to filled water supply tank to reduced foaming (Or) as advised by Veterinarian.

# Piperazine Hydrate 56.3% W/V Oral Solution

## Pipersif-56.3

**Composition:**

Each ml contains  
Piperazine Hydrate I.P 56.3 % w/v  
(Equivalent to Piperazine base 25% w/v)  
Excipients Q.S

**Indications:**

Drug of choice for removal of nematodes and Roundworms (Ascaridia galli, Capillaria Spp.) in poultry

**Dosage:** Orally through drinking water

**Poultry:**

**Chickens:** Mix 60ml Piperazine dewormer solution in 10 liters of water. This will provide medication for 300 birds upto 4-8 weeks of age, mix 60 ml piperazine solution in 13 liters of water for 150 birds over 8-16 weeks of age i.e. 40 ml in 8-9 liters of water for 100 birds (or) as advised by veterinarian.



# Tylosin Tartrate Powder 62.5% w/w

## TYLOSIF

**Composition:**

Each gram contains  
Tylosin Tartrate I.P 625 mg  
Excipients Q.S

**Indications:**

**Poultry:** Prevention and Treatment of CRD, infectious Coryza, Staphylococcosis and as immunity enhancer.

Prevention and control of mycoplasma infections.